

Matroid Matching: the Power of Local Search

[Extended Abstract]

Jon Lee
IBM Research
Yorktown Heights, NY
jonlee@us.ibm.com

Maxim Sviridenko
IBM Research
Yorktown Heights, NY
sviri@us.ibm.com

Jan Vondrák
IBM Research
San Jose, CA
jvondrak@us.ibm.com

ABSTRACT

We consider the classical *matroid matching* problem. Unweighted matroid matching for linear matroids was solved by Lovász, and the problem is known to be intractable for general matroids. We present a PTAS for unweighted matroid matching for general matroids. In contrast, we show that natural LP relaxations have an $\Omega(n)$ integrality gap and moreover, $\Omega(n)$ rounds of the Sherali-Adams hierarchy are necessary to bring the gap down to a constant.

More generally, for any fixed $k \geq 2$ and $\epsilon > 0$, we obtain a $(k/2 + \epsilon)$ -approximation for matroid matching in k -uniform hypergraphs, also known as the matroid k -parity problem. As a consequence, we obtain a $(k/2 + \epsilon)$ -approximation for the problem of finding the maximum-cardinality set in the intersection of k matroids. We have also designed a $3/2$ -approximation for the weighted version of a special case of matroid matching, the *matchoid problem*.

Categories and Subject Descriptors

F.2.2 [Nonnumerical Algorithms and Problems]: Computations on discrete structures

General Terms

Algorithms, Theory

Keywords

matroid, matching, local search, Sherali-Adams hierarchy

1. INTRODUCTION

The matroid matching problem was proposed by Lawler as a common generalization of two important polynomial-time solvable problems: the non-bipartite matching problem, and the matroid-intersection problem (see [32]). Unfortunately, it turns out that matroid matching for general matroids is intractable and requires an exponential number of queries if the matroid is given by an oracle (see [36, 26]). This result

can be easily transformed into an NP-completeness proof for a concrete class of matroids (see [46]). An important result of Lovász is that (unweighted) matroid matching can be solved in polynomial time for *linear matroids* (see [35]). There have been several attempts to generalize Lovász' result to the weighted case. Polynomial-time algorithms are known for some special cases (see [49]), but for general linear matroids there is only a pseudopolynomial-time randomized exact algorithm (see [8, 40]).

In this paper, we revisit the matroid matching problem for general matroids. Our main result is that while LP-based approaches including the Sherali-Adams hierarchy fail to provide any meaningful approximation, a simple local-search algorithm gives a PTAS (in the unweighted case). This is the first PTAS for general matroid matching and to our knowledge also the first example of a problem where there is such a dramatic gap between the performance of the Sherali-Adams hierarchy and a simple combinatorial algorithm. We also provide approximation results for a generalization of the problem to hypergraphs; more details follow.

We assume familiarity with approximation algorithms (see [51], for example) and matroid algorithmics (see [46], for example). Throughout, we consider maximization problems. A c -approximation algorithm finds in polynomial time a solution of value at least OPT/c . Briefly, for a matroid \mathcal{M} , we denote the ground set of \mathcal{M} by $V = V(\mathcal{M})$, its set of independent sets by $\mathcal{I} = \mathcal{I}(\mathcal{M})$, and its rank function by $r_{\mathcal{M}}$. For a given matroid \mathcal{M} , the associated *matroid constraint* is $S \in \mathcal{I}(\mathcal{M})$ or equivalently $|S| = r_{\mathcal{M}}(S)$.

In the *matroid hypergraph matching problem*, we are given a matroid $\mathcal{M} = (V, \mathcal{I})$ and a hypergraph $G = (V, \mathcal{E})$ where $\mathcal{E} \subseteq 2^V$. Note that the vertex set of the hypergraph G and the ground set of the matroid \mathcal{M} are the same. The goal is to choose a maximum-cardinality collection of disjoint hyperedges $E^* \subseteq \mathcal{E}$ in hypergraph G , such that the set of vertices covered by hyperedges in E^* is an independent set in matroid \mathcal{M} . If G is a graph, we obtain the classical matroid matching problem.

The matroid hypergraph matching problem generalizes several classical optimization problems, namely:

1. If \mathcal{M} is a *free matroid* (i.e., $\mathcal{I}(\mathcal{M}) = 2^V$), then the problem is the *maximum hypergraph matching problem* or the *maximum set-packing problem*. Set packing in general is NP-hard, but when G is a graph, it is the classical matching problem which led Edmonds to the notion of polynomial-time algorithms (see [15, 16]).
2. In the *k -matroid intersection problem* we are given k matroids $\mathcal{M}_1 = (V, \mathcal{I}_1), \dots, \mathcal{M}_k = (V, \mathcal{I}_k)$ on the same

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.


STOC 2010, Cambridge, MA
Copyright 2010 ACM ...\$5.00.

ground set V , and the goal is to find a maximum cardinality set S of elements that is independent in each of the k matroids, i.e. $S \in \cap_{j=1}^k \mathcal{I}_j$. The k -matroid intersection problem is NP-hard for $k \geq 3$ but polynomially solvable for $k = 2$ (see [46]).

3. A problem of intermediate generality is the k -uniform matchoid problem, defined for $k = 2$ by Edmonds and studied by Jenkyns (see [27]). In this problem, we have a k -uniform hypergraph and a matroid \mathcal{M}_v given for each vertex v , having ground set the set of hyperedges containing v . The goal is to choose a maximum collection of hyperedges S , such that for each v , the hyperedges in S containing v form an independent set in \mathcal{M}_v . This can be also seen as a packing problem with many matroid constraints, where each item participates in at most k of them.

By taking each \mathcal{M}_v to be the uniform matroid of rank 1, we get the set-packing problem. By taking k arbitrary matroids defined on k copies of the same ground set V and a hypergraph of n parallel hyperedges on the k copies of the same element from V , we get k -matroid intersection. On the other hand, the matchoid problem is a special case of matroid matching, as we show below. We remark that even for $k = 2$, the matchoid problem is NP-hard (see [36]).

4. The special case of the matroid hypergraph matching problem when each vertex belongs to a unique hyperedge, and all hyperedges have cardinality exactly k is known as the *matroid k -parity problem*, or simply the *matroid parity problem* when $k = 2$. As we show below, this problem is in fact equivalent to k -uniform matroid matching, even in terms of approximation.



Next, we explain how the k -uniform matchoid problem is a special case of matroid k -parity. Given a hypergraph G , we can replace each vertex by n_v distinct copies, where n_v is the number of hyperedges containing v . We replace each hyperedge in G by a collection of distinct copies of its elements, so that we get a hypergraph G' where the hyperedges are disjoint. In the matchoid problem, we have a matroid \mathcal{M}_v defined on the n_v copies of each vertex v , and we define a new matroid \mathcal{M}' by taking the union of the matroids \mathcal{M}_v . Then matroid k -parity for (G', \mathcal{M}') is equivalent to the original k -uniform matchoid for (G, \mathcal{M}_v) .

In fact, a similar construction implies that matroid k -parity includes (and therefore is equivalent to) matroid matching in k -uniform hypergraphs (or more generally in hypergraphs where each hyperedge has cardinality at most k , which can be shown by adding dummy elements). Given an instance of k -uniform matroid matching, we define n_v copies for each vertex v where n_v is the degree of v in the hypergraph $G = (V, \mathcal{E})$. We replace each hyperedge in G by a collection of distinct copies of its elements, so that the new hyperedges are disjoint. Let $G' = (V', \mathcal{E}')$ be the new hypergraph. We also define a new matroid \mathcal{M}' on the ground set V' , where the n_v copies of each vertex act as parallel copies. That is, a set of vertices $S' \subseteq V'$ is independent in \mathcal{M}' if it contains at most one copy of each vertex from V and the respective set $S \subseteq V$ is independent in \mathcal{M} . It is not difficult to show that \mathcal{M}' is a matroid.

Henceforth, we will discuss the matroid k -parity problem with the understanding that all our results can be easily extended to the k -uniform matroid matching problem. For the purposes of this paper, the terms *matroid matching* and *matroid parity* are essentially interchangeable.

Literature overview.

There are a few different lines of research relevant to our results. The matroid parity problem (for $k = 2$) was originally popularized by Lawler [31]. The maximum-cardinality matroid parity problem was shown to have exponential query complexity in general (see [36, 26]), and to be NP-hard for some concrete classes of matroids. If the matroid is linear over the reals, the problem is polynomially solvable (see [35, 20, 41, 42, 48]). For more on matroid parity, applications and closely related problems, see [6, 37, 43, 44]. The linear matroid k -parity problem also admits a polynomial time algorithm if k is a constant, and the rank of matroid \mathcal{M} is $O(\log |V|)$ (see [4]). The special case of matroid intersection can be solved in polynomial time for arbitrary matroids (see for example [46]), even in the weighted case.

It is not difficult to show that a simple greedy algorithm that adds hyperedges one-by-one, as long as the current solution remains independent, gives a k -approximation for general weighted matroid k -parity and that this guarantee is tight. This follows from the work of Jenkyns on k -independence systems; see [28] (see also Section 2). Until our present work, this was the only algorithm known for the general matroid k -parity problem with a provable approximation guarantee. The only improvement to our knowledge has been achieved in the case of unweighted matroid matching ($k = 2$), where Fujito proved that a local-search algorithm gives a $3/2$ -approximation (see [19]).

Linear-programming relaxations for the matroid matching problem ($k = 2$) have been studied in [50, 11, 12, 21]. An LP proposed by Vande Vate (see [50]) has been shown to be half-integral, and moreover, there are polynomial-time algorithms to find a half-integral optimal solution in the unweighted case (see [11, 12]) and the general weighted case (see [21]). However, this approach has not yielded any approximation algorithms for matroid matching.

We now survey known results on the approximability of various special cases of the matroid k -parity problem. Local-search algorithms exploring a larger neighborhood for the k -set packing problem were analyzed in [25]. They showed that a local-search algorithm produces a solution with approximation guarantee $k/2 + \epsilon$ for any fixed $\epsilon > 0$ and constant k

in polynomial time. Hardness of approximation results are known for the maximum k -dimensional matching problem which is a special case of the maximum k -set packing problem and hence also matroid k -parity. The best known lower bound is the $\Omega(k/\log k)$ hardness of approximation of [24]. It is also known that a large-neighborhood local-search algorithm has a tight approximation guarantee of $k - 1 + \epsilon$ for the problem of (weighted) k -matroid intersection (see [33]). It should be noted though that unweighted versions of packing problems seem to be easier for algorithm design and analysis, and approximation guarantees for general linear objective functions can often be improved for an unweighted variants.

The Sherali-Adams hierarchy (see [47]) has been studied recently for a number of combinatorial optimization problems (see [29, 30] for surveys on the results in this area). Mathieu and Sinclair proved (see [38]) for the non-bipartite matching problem that r rounds of Sherali-Adams applied to the matching polytope have integrality gap $1 + O(1/r)$ and hence provides a PTAS (while the problem can be solved exactly in polynomial time). Chan and Lau considered k -uniform hypergraph matching, i.e. k -set packing, and proved that that even after $O(n)$ rounds of Sherali-Adams, the standard LP has integrality gap at least $k - 2$ (see [9]). In contrast, local search techniques yield a $(k/2 + \epsilon)$ -approximation (see [25]), and an alternative LP (using intuition from local search) has integrality gap at most $(k + 1)/2$ (see [9]).

Our Results.

On the negative side, we show that the known linear relaxations of matroid parity do not yield any reasonable approximation guarantee (even for $k = 2$ with unit weights). More precisely, all variants of the LPs that have been proposed have an $\Omega(n)$ integrality gap for instances with n pairs. Moreover, $\Omega(n)$ rounds of the Sherali-Adams hierarchy are required to generate an LP with a constant integrality gap.

In contrast, we prove that a very simple local-search algorithm gives a PTAS for unweighted matroid parity. Given the negative results for the matroid parity problem (see [36, 26]), this is the best type of worst-case result we could expect for this problem. It is also a strong manifestation of the fact that LP-based hierarchies do not always match the performance of combinatorial algorithms (which was, in a weaker sense, shown previously for the matching problem in graphs (see [38]) and hypergraphs (see [9])).

For the more general problem of unweighted matroid k -parity, we present a $(k/2 + \epsilon)$ -approximation, for any fixed $k \geq 2$ and $\epsilon > 0$. As a special case, this subsumes the (unweighted) k -matroid intersection problem for which a k -approximation was known since 1976 (see [28]) and has been only recently improved to $k - 1 + \epsilon$ (see [33]).

The algorithm that we analyze is simple local search that in each iteration seeks to remove $s(\epsilon)$ hyperedges and add $s(\epsilon) + 1$ hyperedges to the current solution in such a way that the new solution defines an independent set in matroid \mathcal{M} . We call this the *s-neighborhood local-search algorithm*. If there is no improvement the algorithm stops and outputs the current local optimum. Our analysis uses an idea from [25] to reduce inductively the instance, and given the performance of the local search on the smaller instance, to derive the guarantee on the original one. But the presence of the matroid independence constraint complicates matters significantly. In particular, to achieve the approximation guarantee of $1 + \epsilon$ for the matroid parity problem we need

to implement the local-search algorithm with $s(\epsilon)$ exponentially large in $1/\epsilon$, while it is well known that $s(\epsilon) = \lceil 1 + 1/\epsilon \rceil$ is enough for the maximum matching problem in graphs (the fixed-size augmenting-path algorithm could be viewed as a local-search algorithm). We do not know if having such a large neighborhood is necessary or if it is just an artifact of our analysis. Surprisingly, for $k \geq 3$, we show that to achieve $k/2 + \epsilon$ approximation, it is enough to run the local-search algorithm with $s(\epsilon)$ polynomially bounded in $1/\epsilon$.

The rest of the paper is organized as follows. In §2, we show that matroid k -parity is a special case of a “ k -independence system” which implies a greedy k -approximation. In §3, we present our PTAS for unweighted matroid parity. In §4, we consider various linear-programming relaxations for the matroid parity problem and present our lower bounds on their integrality gap. In §5, we present a $(k/2 + \epsilon)$ -approximation for matroid k -parity.

We have also developed a $3/2$ -approximation algorithm for the weighted matchoid problem, which is a special case of weighted matroid parity. This result uses an LP relaxation of the matchoid problem and its known half-integrality (see [11, 12, 21, 50] for closely related LPs). We provide an alternative proof that is very simple and intuitive which might be of independent interest. Unfortunately, the space here does not allow us to present these results, so we refer to [34].

2. RELATION TO INDEPENDENCE SYSTEMS

First, we show that the matroid k -parity falls in the framework of k -independence systems (see [28]). Such systems generalize intersections of k matroids, and in fact several definitions of various degrees of generality have been proposed (see also [39, 7]). Jenkyns’ definition is as follows.

DEFINITION 1. *A family of sets $\mathcal{I} \subset 2^V$ is a p -system, if for all $W \subseteq V$,*

$$\max\{|B| : B \subseteq W, B \in \mathcal{I}\} \leq p \cdot \min\{|B| : B \subseteq W, B \in \mathcal{I}\}$$

LEMMA 1. *The independence system corresponding to matroid k -parity is a k -system.*

PROOF. Consider an independent collection of hyperedges $W = \{e_1, \dots, e_\ell\}$, and two bases (i.e., setwise maximal independent subsets) B_1, B_2 of W . Assume toward a contradiction that $|B_2| > k|B_1|$. Let $S_1 = \bigcup\{e : e \in B_1\}$ and $S_2 = \bigcup\{e : e \in B_2\}$; i.e. $|S_i| = k|B_i|$ and both S_1 and S_2 are independent in the matroid \mathcal{M} . By the matroid extension axiom, S_1 can be completed from S_2 to a set $S_1 \cup S'_2$ independent in \mathcal{M} , where $S'_2 \subseteq S_2 \setminus S_1$ and $|S'_2| = |S_2| - |S_1| = k|B_2| - k|B_1|$. Note that S'_2 is not necessarily a union of hyperedges. However, it must contain at least one hyperedge, otherwise $|S'_2| \leq (k - 1)|B_2| < k|B_2| - k|B_1|$. Therefore, there is a hyperedge $e_i \in B_2 \setminus B_1$ that we can add to B_1 which contradicts B_1 being a base of W . \square

The work of [27, 18] for p -systems gives the following results (see also [7]).

THEOREM 1. *The greedy algorithm gives a p -approximation for maximizing a linear function over a p -system. Moreover, the greedy algorithm gives a $(p + 1)$ -approximation for maximizing a monotone submodular function over a p -system.*

COROLLARY 1. *The greedy algorithm gives a k -approximation for matroid k -parity even in the weighted version. Moreover, the greedy algorithm gives a $(k + 1)$ -approximation for maximizing a monotone submodular function over sets feasible for the matroid k -parity problem.*

We regard the greedy k -approximation for matroid k -parity as a “folklore” result and a starting point for further improvements. For unweighted matroid parity ($k = 2$), this has been improved to a factor of $3/2$ by Fujito [19]. For general k , no better approximation was known prior to our work.

3. PTAS FOR MATROID PARITY

Let us start with the case of $k = 2$, i.e. matroid parity. In an instance of matroid parity, we have disjoint pairs, and we look for a maximum-cardinality collection of pairs whose union forms an independent set in a given matroid. We present a PTAS for this problem.

DEFINITION 2. *For feasible solutions A and B of matroid parity, a “local move of size s between A and B ” is a choice of $s - 1$ pairs e_1, \dots, e_{s-1} inside A , and s pairs e'_1, \dots, e'_s inside B , such that $(A \setminus \bigcup_{i=1}^{s-1} e_i) \cup \bigcup_{i=1}^s e'_i$ is again feasible.*

THEOREM 2. *For any $\epsilon > 0$, a local-search algorithm which considers local moves of size up to $s(\epsilon) = 5^{\lceil 1/(2\epsilon) \rceil}$ achieves a $(1 + \epsilon)$ -approximation for the matroid parity problem.*

The same result also holds for matroid matching, by a simple reduction that we outlined in the introduction. The theorem follows immediately from the following characterization of local optima.

LEMMA 2. *Let $t \geq 1$, and A, B feasible solutions to the matroid parity problem such that*

$$|A| < (1 - 1/2t) |B|.$$

Then there is a local move of size 5^{t-1} between A and B .

Assuming that B is an actual optimum and A is a local optimum with respect to local moves of size 5^{t-1} , this implies that A is a $2t/(2t - 1)$ -approximate solution. This means that for any fixed $\epsilon > 0$, we can pick $t = \lceil 1/(2\epsilon) \rceil + 1$ and $s = 5^{t-1}$; the corresponding local-search algorithm achieves a $(1 + \epsilon)$ -approximation for matroid parity.

It remains to prove the lemma. Our proofs uses the standard notion of matroid contraction. For a set $S \subset V(\mathcal{M})$, \mathcal{M}/S (read \mathcal{M} contract S) is the matroid having ground set $V(\mathcal{M}) \setminus S$ and set of independent sets $\{T \subseteq V(\mathcal{M}) \setminus S : T \cup J \in \mathcal{I}(\mathcal{M})\}$, where J is an arbitrary maximal independent subset of S with respect to \mathcal{M} .

PROOF. Let A, B be feasible solutions as above. (We assume for simplicity that A and B are disjoint, otherwise we can contract the intersection, which only decreases the ratio $|A|/|B|$.) Because $|A| < |B|$, there exists $B_0 \subset B$, $|B_0| = |B| - |A|$ such that $A \cup B_0$ is independent in \mathcal{M} . We proceed by induction on t .

Base case: $t = 1$.

For $t = 1$, we have $|A| < \frac{1}{2}|B|$. Then, $|B_0| = |B| - |A| > \frac{1}{2}|B|$. Because B decomposes into disjoint pairs, this means there must be a pair contained inside B_0 . This pair can be added to A without violating independence, i.e. there is a local move of size one.

General case: $t \geq 2$.

We assume that $|A| = |B| - a$ where $a > \frac{1}{2t}|B|$. We also assume $a \leq \frac{1}{2}|B|$, otherwise we are in the base case. We construct a set $B_0 \subset B$ as above, with $A \cup B_0$ independent and $|B_0| = a$. Again, if there is a pair contained inside B_0 , we can add it to A , and we are done. So let us assume that no pair is contained completely inside B_0 .

Every pair intersecting B_0 also contains an element in $B \setminus B_0$; let us denote the elements matched with B_0 by B_1 . We have $|B_1| = |B_0| = a$. Let $\mathcal{M}_0 = \mathcal{M}/B_0$ denote the matroid where B_0 has been contracted. Because $A \cup B_0$ and $B_1 \cup B_0$ are independent in \mathcal{M} (by construction), we get that A and B_1 are independent in \mathcal{M}_0 . Because $|A| = |B| - a \geq a = |B_1|$, we can extend B_1 by adding (possibly zero) elements from A , to form an \mathcal{M}_0 -independent set $(A \setminus A_1) \cup B_1$ where $|A_1| = |B_1| = a$.

If A_1 contains a pair e then we can find a local move as follows: $A \setminus e$ is independent in \mathcal{M}_0 , and so is the set $(A \setminus A_1) \cup B_1$. Therefore, $A \setminus e$ can be extended to a set $(A \setminus e) \cup \{x', x''\}$ independent in \mathcal{M}_0 , such that $x', x'' \in B_1$. The elements x', x'' are contained in pairs e', e'' whose remaining elements are in B_0 . Because $(A \setminus e) \cup \{x', x''\}$ is independent in $\mathcal{M}_0 = \mathcal{M}/B_0$, any elements of B_0 can be added for free, and $(A \setminus e) \cup e' \cup e''$ is independent in \mathcal{M} . This defines a local move of size two.

The rest of the proof deals with the case when there is no pair contained in A_1 . Then, every pair intersecting A_1 also contains an element in $A \setminus A_1$; let us denote the elements matched with A_1 by A_2 . We have $|A_2| = |A_1| = a$. Here is where we apply the inductive hypothesis.

The inductive step.

We define a new matroid $\mathcal{M}_1 = \mathcal{M}_0/B_1 = \mathcal{M}/(B_0 \cup B_1)$. By construction, the sets $A^* = A \setminus (A_1 \cup A_2)$ and $B^* = B \setminus (B_0 \cup B_1)$ are both independent in \mathcal{M}_1 . They both form a union of pairs and hence are feasible solutions to the matroid parity problem for \mathcal{M}_1 . We have $|A^*| = |A| - 2a$ and $|B^*| = |B| - 2a$. Because $|A| = |B| - a$, we get

$$\frac{|A^*|}{|B^*|} = \frac{|A| - 2a}{|B| - 2a} = \frac{|B| - 3a}{|B| - 2a} = 1 - \frac{1}{|B|/a - 2}.$$

Because we assumed $a > \frac{1}{2t}|B|$, we have $|B|/a < 2t$ and $|A^*| < (1 - \frac{1}{2t-2})|B^*|$, so we can apply the inductive hypothesis. There is a local move of size $s = 5^{t-2}$ between A^* and B^* , i.e. a union of $s - 1$ pairs $\tilde{A} \subseteq A^*$ and s pairs $\tilde{B} \subseteq B^*$ such that $(A^* \setminus \tilde{A}) \cup \tilde{B}$ is independent in \mathcal{M}_1 . Our goal is to find a local move of size $5s = 5^{t-1}$ between A and B (in \mathcal{M}).

The set $(A^* \setminus \tilde{A}) \cup \tilde{B}$ is independent in \mathcal{M}_1 . Unfortunately, $(A \setminus \tilde{A}) \cup \tilde{B}$ is not necessarily independent, even in \mathcal{M} . We have to proceed more carefully. The set $(A^* \setminus \tilde{A}) \cup A_2 = A \setminus (A_1 \cup \tilde{A})$ is independent in $\mathcal{M}_1 = \mathcal{M}_0/B_1$, because $(A \setminus A_1) \cup B_1$ was constructed to be independent in \mathcal{M}_0 . Therefore, we can extend $(A^* \setminus \tilde{A}) \cup \tilde{B}$ to a set $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ independent in \mathcal{M}_1 , where $C_2 \subseteq A_2$ and $|C_2| \leq |\tilde{B}|$. (If $|A_2| \leq |\tilde{B}|$, we can just set $C_2 = A_2$.)

The new set $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ is also independent in \mathcal{M}_0 (a weaker condition). So is $(A^* \setminus \tilde{A}) \cup (A_2 \setminus C_2) \cup A_1$, as any subset of A is independent in \mathcal{M}_0 . Therefore, we can extend $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ to a set $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2) \cup (A_1 \setminus C_1)$ in \mathcal{M}_0 , where $C_1 \subseteq A_1$ and $|C_1| \leq |\tilde{B}|$.

The set we have obtained is not necessarily a union of pairs, so let us remove the whole pair for each element in C_1 and C_2 . Let us denote by C' the union of all pairs intersecting $C_1 \cup C_2$. By our construction, we have $C_1 \cup C_2 \subseteq C' \subseteq A_1 \cup A_2$. Further, let us define $C'_1 = C' \cap A_1$ and $C'_2 = C' \cap A_2$. Each pair on $A_1 \cup A_2$ contains exactly one element in A_1 and one element in A_2 , therefore $|C'_1| = |C'_2|$. Also, $|C'_1| = |C'_2| \leq |C_1 \cup C_2|$, because each element of $C_1 \cup C_2$ contributes at most one pair to C' .

We obtain a feasible solution $A^+ = (A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C'_2) \cup (A_1 \setminus C'_1)$ in \mathcal{M}_0 . Now, consider the set $(A^+ \setminus A_1) \cup B_1 = (A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C'_2) \cup B_1$. This is independent in \mathcal{M}_0 , because $A^+ \setminus A_1$ was constructed to be independent in $\mathcal{M}_1 = \mathcal{M}_0/B_1$. Because A^+ misses some elements in A_1 , namely C'_1 , the cardinality of $(A^+ \setminus A_1) \cup B_1$ is actually larger than $|A^+|$, $|(A^+ \setminus A_1) \cup B_1| = |A^+| + |C'_1|$. Hence, we can extend A^+ by $|C'_1|$ elements of B_1 , let us call them F_1 , to obtain a set $A^+ \cup F_1$ independent in \mathcal{M}_0 . The pairs touching F_1 have exactly 1 element in B_1 and the other element in B_0 . Let F_0 be the elements of B_0 matched with F_1 . We can add F_0 for free and obtain an independent set $A^+ \cup F_1 \cup F_0$ in \mathcal{M} . We have $|F_0| = |F_1| = |C'_1| = |C'_2|$. Now $A^+ \cup F_1 \cup F_0$ is a union of pairs and hence a feasible solution, of cardinality

$$|A^+ \cup F_1 \cup F_0| = |A^+ \cup C'_1 \cup C'_2| = |(A \setminus \tilde{A}) \cup \tilde{B}| > |A|.$$

Finally, let us estimate the size of this local move. We removed $\tilde{A} \cup C'_1 \cup C'_2$ from A , and added $\tilde{B} \cup F_1 \cup F_0$ instead. The size of C'_1 is bounded by $|C'_1| \leq |C_1 \cup C_2| \leq 2|\tilde{B}|$, hence $|F_1| = |C'_1| \leq 2|\tilde{B}|$. The size of F_0 is equal to the size of F_1 , i.e. $|F_0 \cup F_1| = 2|F_1| \leq 4|\tilde{B}|$. In summary, we are adding at most $5|\tilde{B}|$ elements to A , i.e. the size of the local move is at most $5|\tilde{B}| = 5s = 5^{t-1}$. \square

4. LP RELAXATIONS

In this section, we consider a linear-programming approach to matroid parity. Our results in this direction are mostly negative and indicate that linear programming in this case failed very badly compared to the local-search algorithm presented in the previous section. We formulate our linear programs for the general case of matroid k -parity but the case $k = 2$ is already sufficiently general to obtain our results.

We start with the following natural LP for the weighted matroid k -parity problem (equivalent to an LP studied in [21]). The variables y_e correspond to the hyperedges and the variables x_u correspond to the elements of the ground set. We assume here that each hyperedge alone is independent; otherwise we remove it from the instance.

$$\max \sum_{e \in \mathcal{E}} w_e y_e, \quad (1)$$

$$\sum_{u \in S} x_u \leq r_{\mathcal{M}}(S), \quad \forall S \subseteq V, \quad (2)$$

$$x_u = y_e, \quad \forall u \in e, e \in \mathcal{E}, \quad (3)$$

$$x_u, y_e \geq 0, \quad \forall u \in V, e \in \mathcal{E}. \quad (4)$$

In the objective function (1) we are maximizing the total weight of chosen hyperedges. The constraints (3) correspond to the fact that we can choose a hyperedge only if we chose all its vertices. The constraints (2) are the standard rank constraints for the matroid \mathcal{M} and any independent set of vertices must satisfy them.

For any set of elements $S \subseteq V$ let $\text{sp}(S) = \{u \in V \mid r_{\mathcal{M}}(S \cup \{u\}) = r_{\mathcal{M}}(S)\}$ be the *span* of S in matroid \mathcal{M} . $F \subseteq V$ is a *flat* if $\text{sp}(F) = F$ (for more information on these concepts, see [46], Chapter 39). Since $r_{\mathcal{M}}(\text{sp}(S)) = r_{\mathcal{M}}(S)$, it is easy to see that it is enough to write the constraints (2) for every flat $F \subseteq V$; the inequality for arbitrary $S \subseteq V$ is implied by the flat $F = \text{sp}(S)$.

Another set of valid inequalities (for $k = 2$) were suggested by Vande Vate [50] and studied in subsequent work [11, 12, 21, 50]. For a set $S \subseteq V$ and a hyperedge e , let $a(S, e) = r_{\mathcal{M}}(S \cap \text{sp}(e))$. In case of a flat F , the intuition is that $a(F, e)$ is the dimension of the subspace of F generated by e . The LP proposed by Vande Vate is as follows.

$$\max \sum_{e \in \mathcal{E}} w_e y_e, \quad (5)$$

$$\sum_{e \in \mathcal{E}} a(S, e) y_e \leq r_{\mathcal{M}}(S), \quad \forall S \subseteq V, \quad (6)$$

$$y_e \geq 0, \quad \forall e \in \mathcal{E}. \quad (7)$$

Again, it is equivalent to consider the inequalities (6) only for flats, which was the formulation given by Vande Vate. This LP is potentially stronger than LP (1-4), which can be equivalently obtained from (5-7) by replacing $a(S, e)$ with the smaller quantity $|S \cap e|$.

It is known that the linear program (5-7) is half-integral in the 2-uniform case. Moreover, there are polynomial time algorithms to find a half-integral optimal solution in the unweighted 2-uniform case (see [11, 12]) and the weighted 2-uniform case (see [21]). The following lemma shows the validity of the LP (5-7) in the general k -uniform case. We could not find a published proof of validity (even for $k = 2$), so for completeness we provide a short proof.

LEMMA 3. *The inequalities (6) are valid for the matroid k -parity problem.*

PROOF. Consider any feasible solution, a collection of hyperedges $E^* = \{e_1, \dots, e_k\}$ such that $e_1 \cup \dots \cup e_k$ is an independent set in matroid \mathcal{M} . In the following, we denote the rank function of \mathcal{M} simply by $r(S)$. Let $S_i = S \cap \text{sp}(e_i)$. Note that $r(S_i) = a(S, e_i)$. We claim that for any $i < k$, $r(S_1 \cup \dots \cup S_i) = r(S_1 \cup \dots \cup S_{i-1}) + r(S_i)$. By induction, we will get that $r(S_1 \cup \dots \cup S_k) = \sum_{i=1}^k r(S_i) = \sum_{i=1}^k a(S, e_i)$ which implies the inequalities (6).

We let $r_A(S) = r(A \cup S) - r(A)$; due to submodularity, this is a non-increasing function of A . Let $A = S_1 \cup \dots \cup S_{i-1}$ and $B = e_1 \cup \dots \cup e_{i-1}$. Our goal is to prove that $r_A(S_i) = r(S_i)$. Because $A \subseteq \text{sp}(B)$, we get $r_A(S_i) \geq r_{\text{sp}(B)}(S_i) = r_B(S_i)$, using that $r(\text{sp}(B)) = r(B)$ and $r(S_i \cup \text{sp}(B)) = r(S_i \cup B)$. On the other hand, as e_i is independent of B , we have

$$\begin{aligned} r(\text{sp}(e_i)) = r_B(\text{sp}(e_i)) &= r_B(S_i) + r_{B \cup S_i}(\text{sp}(e_i)) \\ &\leq r_A(S_i) + r_{S_i}(\text{sp}(e_i)) \end{aligned}$$

using again the submodularity of r . This implies that $r_A(S_i) \geq r(\text{sp}(e_i)) - r_{S_i}(\text{sp}(e_i)) = r(S_i)$. The opposite inequality is obvious and hence $r_A(S_i) = r(S_i)$. \square

In the following, we use examples where $e = \text{sp}(e)$ for all hyperedges $e \in \mathcal{E}$. Note that in this case, $a(S, e) = r_{\mathcal{M}}(S \cap \text{sp}(e)) = r_{\mathcal{M}}(S \cap e) = |S \cap e|$, and hence the two LPs are in fact equivalent.

4.1 Integrality gap example

It is known that the integrality gap of the LP relaxation (1–4) is $k - 1 + \frac{1}{k}$ for the maximum weighted hypergraph matching problem [9]. Therefore, it is tempting to conjecture that a similar result should hold for matroid hypergraph matching. Unfortunately, as we show below, the integrality gap of the linear-programming relaxation (1–4) is $\Omega(|\mathcal{E}|)$ even when $k = 2$ and the matroid is linear over the rationals.

Example.

Consider a ground set $V = \{u_1, v_1, \dots, u_n, v_n\}$ of size $2n$, partitioned into pairs $e_i = \{u_i, v_i\}$. The weight of each pair is $w_{e_i} = 1$. Given an integer parameter $t \geq 1$, we define a matroid $\mathcal{M} = (V, \mathcal{I})$ as follows. For a set $S \subseteq V$, let $p(S)$ be the number of pairs e_i such that $e_i \subseteq S$. Then let $S \in \mathcal{I}$ if $p(S) \leq t$.

It can be checked that \mathcal{I} satisfies the matroid independence axioms: For any $S, T \in \mathcal{I}$, $|S| < |T|$, either T contains an element from a pair $\{u_i, v_i\}$ which is disjoint from S , or it contains more pairs than S . In either case, we can extend S by adding some element of T . Moreover, this matroid is linear over the rationals (see [34]).

First, let us write down the LP for this particular example. We have variables y_i for $i = 1, \dots, n$, which are constrained by $y_i \in [0, 1]$. Since $\text{sp}(e_i) = e_i$, the LPs (1–4) and (5–7) coincide. It is enough to write the constraints (6) for flats, and in particular only for collections of pairs $S = \bigcup_{i \in T} e_i$. This is because including only one element of a pair in S always increases $r_{\mathcal{M}}(S)$ by 1 and hence cannot strengthen the constraint. Also, for $S = \bigcup_{i \in T} e_i$ where $|T| \leq t$, the rank is $r_{\mathcal{M}}(S) = |S|$ and the respective constraint (6) is implied by $y_i \leq 1$. The only non-trivial constraints are for $S = \bigcup_{i \in T} e_i$, $|T| > t$, where we get $r_{\mathcal{M}}(S) = 2t + (|T| - t) = t + |T|$. Also, $a(S, e_i) = 2$ for all $i \in T$. Therefore, the LP is as follows.

$$\max \sum_{i=1}^n w_i y_i, \quad (8)$$

$$\sum_{i \in T} y_i \leq \frac{1}{2}(t + |T|), \quad \forall T \subseteq [n], |T| > t \quad (9)$$

$$0 \leq y_i \leq 1, \quad \forall i. \quad (10)$$

LEMMA 4. *The integrality gap of LP (8–10) is $\Omega(n/t)$, even in the unweighted case.*

PROOF. It is easy to see that $y_i = 1/2$ for all $i = 1, \dots, n$ is a feasible fractional solution. Therefore, $LP \geq n/2$. However, only t pairs can be selected in an integral optimum, i.e. $OPT = t$. \square

For $t = 1$, we get an $\Omega(n)$ integrality gap. One way to improve the quality of linear-programming relaxations is to add valid inequalities that cut bad fractional solutions. One of the possible classes of valid inequalities are the so-called *clique inequalities* that were recently shown to reduce the integrality gap for unweighted hypergraph matching from $k - 1$ to $(k + 1)/2$ [9]. This motivates us to define the undirected graph $G' = (\mathcal{E}, E')$ where the vertices are the hyperedges $e \in \mathcal{E}$ in our instance of matroid hypergraph matching and the edges are defined between “incompatible hyperedges” e and e' , i.e. when $r(e \cup e') < |e \cup e'|$. A set of vertices C in graph G' is called a clique if it has an edge between every

pair of vertices in C . Let \mathcal{C} be the set of all cliques in graph G' . Then the following set of constraints is valid for the matroid hypergraph matching problem

$$\sum_{e \in C} y_e \leq 1, \quad \forall C \in \mathcal{C}. \quad (11)$$

However, as we can see in the example above (for $t \geq 2$), sometimes the clique inequalities do not add any non-trivial constraints and the LP effectively remains the same. More generally, we could add all the valid constraints for the stable-set polytope corresponding to G' (or perhaps consider the semidefinite program corresponding to the Lovász θ -function). The relaxation would still remain the same, since the graph G' is empty in our example.

In the next section, we consider the strongest known systematic way of generating valid constraints in linear programming, which is the *Sherali-Adams hierarchy*.

4.2 The Sherali-Adams hierarchy

The Sherali-Adams hierarchy produces progressively stronger refinements of a given LP by introducing new variables y_L indexed by subsets of the original variables, and then projecting back to the space of the original variables. We follow the formalism of [38]. To carry out r rounds of Sherali-Adams, we consider all pairs of disjoint subsets of variables I, J such that $|I \cup J| = r$. We multiply each constraint by $\prod_{i \in I} y_i \prod_{j \in J} (1 - y_j)$, expand all the monomial terms and replace every square y_i^2 by y_i . Now all terms are multilinear, and we replace each occurrence of $\prod_{\ell \in L} y_\ell$ by a new variable y_L . We also do the same for the constraint $\prod_{i \in I} y_i \prod_{j \in J} (1 - y_j) \geq 0$, for all disjoint I, J such that $|I \cup J| = r + 1$. This defines the new LP; note that the variables y_L for $|L| > 1$ play no role in the objective function and thus the polytope can be viewed as projected back to the original space.

LEMMA 5. *The integrality gap of LP (8–10) still remains $\Omega(n/r)$ after r rounds of the Sherali-Adams hierarchy.*

PROOF. Our starting point is LP (8–10), with the parameter t chosen equal to the desired number of rounds r . We have constraints $\sum_{\ell \in T} y_\ell \leq \frac{1}{2}(r + |T|)$ for all $|T| > r$. We multiply this constraint by $\prod_{i \in I} y_i \prod_{j \in J} (1 - y_j)$ and obtain

$$\sum_{\ell \in T} y_\ell \prod_{i \in I} y_i \prod_{j \in J} (1 - y_j) \leq \frac{1}{2}(r + |T|) \prod_{i \in I} y_i \prod_{j \in J} (1 - y_j). \quad (12)$$

We expand the products, linearize the expressions, and replace monomials $\prod_{\ell \in L} y_\ell$ by new variables y_L as explained above. We also do the same thing for the constraints $\prod_{i \in I} y_i \prod_{j \in J} (1 - y_j) \geq 0$ with $|I \cup J| = r + 1$. We claim that $y_L = 1/2^{|L|}$ for all $|L| \leq r + 1$ is a feasible solution for the new LP.

To see this, first observe that whenever we have $\ell \in J$ on the left-hand side of (12), the corresponding term contains $y_\ell(1 - y_\ell) = y_\ell - y_\ell^2$ which disappears after linearization. In terms where $\ell \in I$, we get y_ℓ^2 which gets linearized to y_ℓ . Equivalently, we can replace y_ℓ by 1 in its appearance before the product $\prod_{i \in I} y_i$, whenever $\ell \in I$. Variables outside of $I \cup J$ remain unchanged. Therefore, after linearization, the left-hand side is equal to $(|T \cap I| + \sum_{\ell \in T \setminus (I \cup J)} y_\ell) \prod_{i \in I} y_i \prod_{j \in J} (1 - y_j)$.

Now we replace the monomials $\prod_{\ell \in L} y_\ell$ by y_L and substitute $y_L = 1/2^{|L|}$. Note that this is equivalent to directly

substituting $y_\ell = 1/2$ for all ℓ . Thus the left-hand side becomes

$$\begin{aligned} & (|T \cap I| + \frac{1}{2}|T \setminus (I \cup J)|) 2^{-|I \cup J|} \\ &= \frac{1}{2}(|T \cap I| + |T \setminus J|) 2^{-|I \cup J|} \leq \frac{1}{2}(r + |T|) 2^{-|I \cup J|} \end{aligned}$$

using the fact that $|I| \leq r$. This verifies the linearized form of constraint (12).

The inequalities arising from $\prod_{i \in I} y_i \prod_{j \in J} (1 - y_j) \geq 0$ are easy to verify, since our assignment $y_L = 1/2^{|L|}$ is equivalent to substituting $y_i = 1/2$. Therefore, our fractional solution is feasible for r rounds of Sherali-Adams.

Finally, the value of our fractional solution is equal to $n/2$, because each singleton variable is $y_i = y_{\{i\}} = 1/2$. The integral optimum is $OPT = r$. \square

To summarize, our LP (8–10) is an instance of the strongest “natural LP” for matroid matching we are aware of, namely the Vande Vate LP (5–7). The same LP (8–10) is obtained even with the added clique constraints (11) and other valid constraints for the stable-set polytope which are hard to optimize over in general. On top of this LP, we run the Sherali-Adams hierarchy and the gap still remains superconstant for $o(n)$ rounds.

4.3 Lower bound on Chvátal rank

Another popular way to derive progressively stronger linear-programming relaxations is to apply Chvátal-Gomory cuts (for example, see [45]). Let $P := \{x \in \mathbb{R}^n : \sum_{j=1}^n a_{i,j} x_j \leq b_i, \text{ for } i = 1, \dots, m\}$ be a polyhedron defined by rational data. Let $\sum_j a_j x_j \leq b$ be a linear inequality satisfied by all points in P . Such an inequality is called a *valid (linear) inequality* for P .

Now, let $P_I := \text{conv}(P \cap \mathbb{Z}^n)$. If $\sum_j a_j x_j \leq b$ is a valid linear inequality for P , and $a_j \in \mathbb{Z}$ for all j , then clearly $\sum_j a_j x_j \leq \lfloor b \rfloor$ is a valid linear inequality for P_I . Any such inequality is known as a *Chvátal-Gomory cut* with respect to P . Applying all Chvátal-Gomory cuts to P , we obtain the *first Chvátal closure* $P^{(1)}$ of P . It is a classical result that $P^{(1)}$ is again a polyhedron; that is, only a finite number of the Chvátal-Gomory cuts for P are needed to describe $P^{(1)}$. Now, we can repeatedly apply this closure operator, and we obtain after r repetitions the r -th *Chvátal closure* $P^{(r)}$ of P . If the linear inequality $\sum_j a_j x_j \leq b$ is valid for $P^{(r)}$ but not $P^{(r-1)}$, then we say that the inequality has *Chvátal rank* r . It is a classical result that when P is described by rational data, $P_I = P^{(r)}$ for some finite r .

Hartmann established lower bounds on the *Chvátal rank* for many classes of polytopes of interest in combinatorial optimization (see [22] and also [14]). Eisenbrand and Schulz established that for polytopes in $[0, 1]^n$, the *Chvátal rank* is at most $3n^2 \log(n)$, and furthermore that there is a family of polytopes in the $[0, 1]^n$ that has *Chvátal rank* at least $(1 + \epsilon)n$ (see [17]).

The proof of the following lemma is a generalization of the geometric proof from [14] of the classical result by Chvátal [13] that a clique inequality involving n variables requires at least $\lceil \log_2 n \rceil$ rounds of the Chvátal-Gomory hierarchy. Let \mathcal{P}_t be the polytope described by (8–10). So $\mathcal{P}_t^{(r)}$ is its r -th Chvátal closure.

LEMMA 6. *The point $y^r \in \mathbb{R}^n$ defined by $y_i^r = \frac{1}{2}(\frac{t}{t+1})^r$, for $i = 1, \dots, n$, is in $\mathcal{P}_t^{(r)}$.*

PROOF. We prove the lemma by induction on the number r of rounds of the Chvátal closure. The base case $r = 0$ is trivial since the solution $(1/2, \dots, 1/2)$ is obviously feasible for the linear program (8–10) for any parameter $t \geq 1$.

We assume that $y^{r-1} \in \mathcal{P}_t^{(r-1)}$. Let $\sum_{i=1}^n a_i y_i \leq b$ be a valid linear inequality for $\mathcal{P}_t^{(r-1)}$, with $a_i \in \mathbb{Z}$. Because $(0, \dots, 0)$ is a feasible integral solution for the linear program (8–10), we obtain $(0, \dots, 0) \in \mathcal{P}_t^{(r-1)}$, and therefore $b \geq 0$. If $a_i \leq 0$ for all $i = 1, \dots, n$, then $\sum_{i=1}^n a_i y_i^r \leq 0 \leq \lfloor b \rfloor \leq b$.

Let $A^+ := \sum_{i|a_i > 0} a_i$. If the vector (a_1, \dots, a_n) has at most t strictly positive components, then $A^+ \leq b$ since any integral solution with at most t variables equal to one and the remaining variables equal to zero is a feasible integral solution for the linear program (8–10), which implies that such a solution must belong to $\mathcal{P}_t^{(r-1)}$. Moreover, because A^+ is an integer, we have $A^+ \leq \lfloor b \rfloor$. Therefore, we have $\sum_{i=1}^n a_i y_i^r < A^+ \leq \lfloor b \rfloor$.

Consider now an inequality in which the vector (a_1, \dots, a_n) has more than t strictly positive components. Recall that $a_i \in \mathbb{Z}$ and hence these coefficients are at least 1. An integral solution χ having any t of the respective coordinates equal to one and remaining coordinates equal to zero is feasible for (8–10); therefore $t \leq \sum_{i=1}^n a_i \chi_i \leq b$. We obtain

$$\sum_{i=1}^n a_i y_i^r = \frac{t}{t+1} \sum_{i=1}^n a_i y_i^{r-1} \leq \frac{t}{t+1} b \leq \lfloor b \rfloor,$$

which implies that $y^r \in \mathcal{P}^{(r)}$. \square

COROLLARY 2. *The integrality gap of the linear-programming relaxation obtained from the linear-programming relaxation (8–10) after t rounds of Chvátal closure is $\Omega(n/t)$.*

PROOF. By the Lemma 6 there exists a feasible fractional solution of value $\frac{n}{2}(\frac{t}{t+1})^t = \Omega(n)$ in $\mathcal{P}_t^{(t)}$, while the value of the integral optimal solution is t . \square

COROLLARY 3. *The Chvátal rank of the polytope defined by (8–10) with $t = n/4$ is at least $n/4$.*

PROOF. Consider the inequality $\sum_{i=1}^n y_i \leq t$. This inequality is valid for all integer feasible solutions of the linear programming relaxation (8–10). Let r be the smallest integer such that this inequality is valid for $\mathcal{P}_t^{(r)}$; i.e., r is the Chvátal rank of the inequality $\sum_{i=1}^n y_i \leq t$. By Lemma 6, we obtain $\frac{n}{2}(\frac{t}{t+1})^r \leq t$. Therefore,

$$r \geq \frac{\log_2 n - \log_2(2t)}{\log_2(1 + 1/t)} \geq t(\log_2 n - \log_2(2t)).$$

For $t = n/4$, the Chvátal rank is $r \geq n/4$. \square

5. MATROID K -PARITY

Here we extend the analysis of local search to matroid k -parity; i.e. instead of pairs, we work with hyperedges of size k . We assume that $k \geq 3$. In an instance of matroid k -parity, all hyperedges are mutually disjoint. We remark that our analysis extends to k -uniform matroid matching where hyperedges need not be disjoint, by a standard reduction.

Interestingly, the analysis for $k \geq 3$ is slightly different and the complexity of our $(k/2 + \epsilon)$ -approximation for $k \geq 3$ has a much better dependence on ϵ than our PTAS for $k = 2$ (matroid matching). More precisely, while we need local moves of size exponential in $1/\epsilon$ in order to achieve a

$(1 + \epsilon)$ -approximation for matroid matching, local moves of size polynomial in $1/\epsilon$ are sufficient to achieve a $1/(2/k - \epsilon)$ -approximation for matroid k -parity. We do not know whether our analysis is optimal in terms of this dependence.

DEFINITION 3. For feasible solutions A, B of matroid k -parity, a local move of size s between A and B is a choice of $s - 1$ hyperedges e_1, \dots, e_{s-1} from A , and s hyperedges e'_1, \dots, e'_s from B , such that $(A \setminus \bigcup_{i=1}^{s-1} e_i) \cup \bigcup_{i=1}^s e'_i$ is feasible.

THEOREM 3. For any $k \geq 3$ and $\epsilon > 0$, a local-search algorithm which considers local moves of size up to $s(\epsilon) = \lceil 1/\epsilon^3 \rceil$ achieves a $1/(2/k - \epsilon)$ -approximation for the matroid k -parity problem.

This follows easily from the following characterization of local optima.

LEMMA 7. Let $k \geq 3$, $t \geq 1$, and A, B feasible solutions to the matroid k -parity problem such that

$$|A| < \left(\frac{2}{k} - \frac{1}{(k-1)^t} \right) |B|.$$

Then there exists a local move of size $(2k+1)^{t-1}$ between A and B .

Note that in order to achieve a $1/(2/k - \epsilon)$ -approximation, it suffices to pick $t = \lceil \log_{k-1}(1/\epsilon) \rceil$ and $s(\epsilon) = (2k+1)^{t-1} \leq 1/\epsilon^{\log_{k-1}(2k+1)}$. Then, if A is a local optimum and B is a global optimum, the lemma implies that $|A| \geq (2/k - 1/(k-1)^t)|B| \geq (2/k - \epsilon)|B|$. For simplicity, we replaced $1/\epsilon^{\log_{k-1}(2k+1)}$ by $1/\epsilon^3$ in the statement of the theorem, but for large k the dependency gets close to $1/\epsilon$.

It remains to prove the lemma.

PROOF. (Lemma 7). Let A, B be feasible solutions as above. (We assume for simplicity that A and B are disjoint, otherwise we can contract the intersection, which only decreases the ratio $|A|/|B|$.) Because $|A| < |B|$, there exists $B' \subset B$, $|B'| = |B| - |A|$ such that $A \cup B'$ is independent in \mathcal{M} . We proceed by induction on t .

Base case: $t = 1$.

Here, we have $|A| < (\frac{2}{k} - \frac{1}{k-1})|B| < \frac{1}{k}|B|$. Then, it is impossible that every hyperedge in B contains some element in $B \setminus B'$, because that would mean that $|A| = |B \setminus B'| \geq \frac{1}{k}|B|$. Hence, there must be a hyperedge contained completely inside B' , which can be added to A without violating independence. This means there is a local move of size one.

General case: $t \geq 2$.

We assume that $|A| = (2/k - \epsilon)|B|$ and $\epsilon > \frac{1}{(k-1)^t}$. Again, if there is a hyperedge contained inside B' , we can add it to A , and we are done. So let us assume that no hyperedge is contained completely inside B' .

We use a counting argument to show that there must be many hyperedges with exactly $k-1$ elements in B' . Let a denote the number of such hyperedges ($|e \cap B'| = k-1$), and b the number of hyperedges such that $|e \cap B'| \leq k-2$. All hyperedges in B fall into one of these two categories, hence $|B| = k(a+b)$. On the other hand, $|B'| \leq (k-1)a + (k-2)b$ which means that $|A| = |B| - |B'| \geq a + 2b$. We assumed that $|A| = (2/k - \epsilon)|B|$, which implies

$$a + 2b \leq |A| = (2/k - \epsilon)|B| = (2 - k\epsilon)(a + b). \quad (13)$$

We conclude that

$$a \geq k\epsilon(a + b). \quad (14)$$

Let Q denote these a hyperedges in B and $V(Q)$ denote the elements of B that belongs to hyperedges in Q ; each of them contains exactly one element in $B \setminus B'$ and $k-1$ elements in B' . Let $B_0 = V(Q) \cap B'$ and $B_1 = V(Q) \cap (B \setminus B')$. We have $|B_0| = (k-1)a$ and $|B_1| = a$.

Let $\mathcal{M}_0 = \mathcal{M}/B_0$ denote the matroid with B_0 contracted. Because $A \cup B_0 \subseteq A \cup B'$ and $B_1 \cup B_0 \subseteq B$, both of which are independent in \mathcal{M} , we get that A and B_1 are independent in \mathcal{M}_0 . Because $|A| \geq a + 2b \geq |B_1|$, we can extend B_1 by adding (possibly zero) elements from A , to form a \mathcal{M}_0 -independent set $(A \setminus A_1) \cup B_1$ where $|A_1| = |B_1| = a$.

If A contains any hyperedge e with $|e \cap A_1| \geq 2$, we find a local move of size two as follows: $((A \setminus e) \setminus A_1) \cup B_1$ is an independent set in \mathcal{M}_0 , whose cardinality is at least $|A \setminus e| + 2$ (because A_1 contains ≥ 2 elements of e). Therefore, $A \setminus e$ can be extended to a set $(A \setminus e) \cup \{x', x''\}$ independent in \mathcal{M}_0 , such that $x', x'' \in B_1$. The elements x', x'' are contained in hyperedges e', e'' whose remaining elements are in B_0 . Because $(A \setminus e) \cup \{x', x''\}$ is independent in $\mathcal{M}_0 = \mathcal{M}/B_0$, any elements of B_0 can be added for free, and $(A \setminus e) \cup e' \cup e''$ is independent in \mathcal{M} . This defines a local move of size two.

The rest of the proof deals with the case in which there is no hyperedge in A with more than 1 element in A_1 . Let P be the collection of hyperedges in A intersecting A_1 ; each such hyperedge satisfies $|e \cap A_1| = 1$, and so $|P| = |A_1| = a$. Let A_2 denote the remaining elements of P , i.e. $A_2 \subseteq A \setminus A_1$ and $|A_2| = (k-1)a$. Here we apply the inductive hypothesis.

The inductive step.

We define a new matroid $\mathcal{M}_1 = \mathcal{M}_0/B_1 = \mathcal{M}/(B_0 \cup B_1)$. By construction, the sets $A^* = A \setminus (A_1 \cup A_2)$ and $B^* = B \setminus (B_0 \cup B_1)$ are both independent in \mathcal{M}_1 . They both form a union of hyperedges and hence feasible solutions to the matroid k -parity problem for \mathcal{M}_1 . We have $|A^*| = |A| - ka$ and $|B^*| = |B| - ka = kb$. Using (13), we get

$$\frac{|A^*|}{|B^*|} = \frac{|A| - ka}{kb} = \frac{(2 - k\epsilon)(a + b) - ka}{kb} = \frac{2}{k} - \epsilon - \frac{(k-2 + k\epsilon)a}{kb}$$

and applying (14) to estimate $a \geq k\epsilon$, we get

$$\frac{|A^*|}{|B^*|} \leq \frac{2}{k} - \epsilon - (k-2 + k\epsilon)\epsilon \leq \frac{2}{k} - (k-1)\epsilon.$$

Because we assumed $\epsilon > \frac{1}{(k-1)^t}$, we have

$$|A^*| < \left(\frac{2}{k} - \frac{1}{(k-1)^{t-1}} \right) |B^*|,$$

and we can apply the inductive hypothesis. There is a local move of size $s = (2k+1)^{t-2}$ between A^* and B^* , i.e. a union of $s-1$ hyperedges $\tilde{A} \subseteq A^*$ and s hyperedges $\tilde{B} \subseteq B^*$ such that $(A^* \setminus \tilde{A}) \cup \tilde{B}$ is independent in \mathcal{M}_1 . Our goal is to find a local move of size $(2k+1)s$ between A and B (in \mathcal{M}).

We accomplish this by a construction essentially identical to the case of matroid parity. The set $(A^* \setminus \tilde{A}) \cup \tilde{B}$ is independent in \mathcal{M}_1 . Unfortunately, $(A \setminus \tilde{A}) \cup \tilde{B}$ is not necessarily independent, even in \mathcal{M} . However, the set $(A^* \setminus \tilde{A}) \cup A_2 = A \setminus (A_1 \cup \tilde{A})$ is independent in $\mathcal{M}_1 = \mathcal{M}_0/B_1$, because $(A \setminus A_1) \cup B_1$ was constructed to be independent in \mathcal{M}_0 . Therefore, we can extend $(A^* \setminus \tilde{A}) \cup \tilde{B}$ to a set

$(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ independent in \mathcal{M}_1 , where $C_2 \subseteq A_2$ and $|C_2| \leq |\tilde{B}|$. (If $|A_2| \leq |\tilde{B}|$, we just take $C_2 = A_2$.)

The new set $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ is also independent in \mathcal{M}_0 (a weaker condition). So is $(A^* \setminus \tilde{A}) \cup (A_2 \setminus C_2) \cup A_1$, as any subset of A is independent in \mathcal{M}_0 . So, we can extend $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2)$ to a set $(A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C_2) \cup (A_1 \setminus C_1)$ independent in \mathcal{M}_0 , where $C_1 \subseteq A_1$ and $|C_1| \leq |\tilde{B}|$.

The set we have obtained is not necessarily a union of hyperedges, so let us remove the entire hyperedge for each element in C_1 and C_2 . Let us denote by C' the union of all hyperedges intersecting $C_1 \cup C_2$. Note that due to our construction, $C_1 \cup C_2 \subseteq C' \subseteq A_1 \cup A_2$. We also define $C'_1 = C' \cap A_1$ and $C'_2 = C' \cap A_2$. We know that each hyperedge on $A_1 \cup A_2$ contains exactly one element in A_1 and $k-1$ elements in A_2 . Therefore, $|C'_2| = (k-1)|C'_1|$, and also $|C'_1| = \frac{1}{k}|C'| \leq |C_1 \cup C_2|$, because each element of $C_1 \cup C_2$ contributes at most one hyperedge to C' .

We obtain a feasible solution $A^+ = (A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C'_2) \cup (A_1 \setminus C'_1)$ in \mathcal{M}_0 . Now, consider the set $(A^+ \setminus A_1) \cup B_1 = (A^* \setminus \tilde{A}) \cup \tilde{B} \cup (A_2 \setminus C'_2) \cup B_1$. This is independent in \mathcal{M}_0 , because $A^+ \setminus A_1$ was constructed to be independent in $\mathcal{M}_1 = \mathcal{M}_0/B_1$. Because A^+ misses some elements in A_1 , namely C'_1 , the cardinality of $(A^+ \setminus A_1) \cup B_1$ is actually larger than $|A^+|$, namely $|(A^+ \setminus A_1) \cup B_1| = |A^+| + |C'_1|$. Hence, we can extend A^+ by $F_1 \subseteq B_1$, $|F_1| = |C'_1|$, to obtain a set $A^+ \cup F_1$ independent in \mathcal{M}_0 . The hyperedges touching F_1 have exactly 1 element in B_1 and the remaining $k-1$ elements in B_0 (denote these by F_0), hence we can add F_0 for free and obtain an independent set $A^+ \cup F_1 \cup F_0$ in \mathcal{M} . We have $|F_1| = |C'_1|$ and $|F_0| = (k-1)|F_1| = (k-1)|C'_1| = |C'_2|$. To conclude, we have found a feasible solution $A^+ \cup F_1 \cup F_0$ in \mathcal{M} , of cardinality

$$|A^+ \cup F_1 \cup F_0| = |A^+ \cup C'_1 \cup C'_2| = |(A \setminus \tilde{A}) \cup \tilde{B}| > |A|.$$

Finally, let us estimate the size of this local move. We removed $\tilde{A} \cup C'_1 \cup C'_2$ from A , and added $\tilde{B} \cup F_0 \cup F_1$ instead. The size of C'_1 is bounded by $|C'_1| \leq |C_1 \cup C_2| \leq 2|\tilde{B}|$, hence $|F_0 \cup F_1| = k|C'_1| \leq 2k|\tilde{B}|$. In summary, we are adding at most $(2k+1)|\tilde{B}|$ elements to A , i.e. the size of the local move is at most $(2k+1)|\tilde{B}| = (2k+1)^{t-1}$. \square

6. CONCLUSION

We have seen that a simple combinatorial algorithm performs dramatically better than any known LP-based approach for matroid matching. Linear programming still holds some promise for the k -uniform matchoid problem. Our $\frac{3}{2}$ -approximation for the weighted matchoid problem and the results from [9] on the integrality gap of the hypergraph matching problem motivate the following.

CONJECTURE 1. *The integrality gap of the LP relaxation (1-4) is $k-1 + \frac{1}{k}$ for the maximum weighted k -matchoid problem and $k-1$ for the maximum weighted k -matroid intersection problem.*

In the case of weighted matroid k -parity, we have the following conjecture, which is true (and tight) for the weighted k -set packing problem due to [1, 3] and also for the weighted k -matroid intersection problem due to [33].

CONJECTURE 2. *The simple local-search algorithm for the weighted matroid k -parity problem that tries to add/remove*

a constant number of hyperedges in each iteration has approximation guarantee $k-1+\varepsilon$ for any $\varepsilon > 0$ (with running time depending on $1/\varepsilon$).

A very intriguing open problem is to show that this simple local-search algorithm gives a PTAS for weighted matroid parity ($k=2$). This problem is interesting even for the special case of linear matroids, because Lovász' polynomial-time algorithm applies only to the unweighted case (see [35]). For the weighted linear case, there is only a pseudopolynomial-time randomized exact algorithm [8] and pseudopolynomial-time randomized parallel exact algorithm [40].

Another interesting line of research is to analyze more sophisticated local-search algorithms (see [5, 10]) implemented for the weighted matroid k -parity problem. Such algorithms are known to provide improved approximation guarantees for the weighted set packing problem.

7. REFERENCES

- [1] E. Arkin and R. Hassin, On local search for weighted k -set packing, *Math. of Op. Res.* 23(3) (1998) 640–648.
- [2] M. Balinski, On maximum matching, minimum covering and their connections, in “Proceedings of Princeton Symp. on Math. Prog.,” Eds. H. Kuhn, 303–312, Princeton Univ. Press, 1970.
- [3] V. Bafna, B. Narayanan and R. Ravi, Nonoverlapping local alignments (weighted independent sets of axis-parallel rectangles), *Discrete Applied Mathematics* 71, (1996), no. 1-3, 41–53.
- [4] A. Barvinok, New algorithms for linear k -matroid intersection and matroid k -parity problems, *Math. Program.* 69, 449–470 (1995).
- [5] P. Berman, A $d/2$ approximation for maximum weight independent set in d -claw free graphs, *Nordic J. Comput.* 7(3) (2000), 178–184.
- [6] P. Berman, M. Fürer and A. Zelikovsky, Applications of the Linear Matroid Parity Algorithm to Approximating Steiner Trees, in “Proceedings of CSR 2006,” *LNCS* v.3967, 70–79.
- [7] G. Calinescu, C. Chekuri, M. Pál and J. Vondrák, Maximizing a submodular set function subject to a matroid constraint, to appear in *SIAM Journal on Computing*, special issue for STOC 2008.
- [8] P. Camerini, G. Galbiati and F. Maffioli, Random pseudo-polynomial algorithms for exact matroid problems, *Journal of Algorithms*, 13:258-273, 1992.
- [9] Y.H. Chan and L.C. Lau, On linear and semidefinite programming relaxations for hypergraph matching, to appear in SODA 2010.
- [10] B. Chandra and M. Halldórsson, Greedy local improvement and weighted set packing approximation, *J. Algorithms* 39(2) (2001), 223–240.
- [11] S. Chang, D. Llewellyn and J. Vande Vate, Matching 2-lattice polyhedra: finding a maximum vector, *Discrete Mathematics* 237 (2001), 29–61.
- [12] S. Chang, D. Llewellyn and J. Vande Vate, Matching 2-lattice polyhedra: duality and extreme points, *Discrete Mathematics* 237 (2001), no. 1–3, 63–95.
- [13] V. Chvátal, Edmonds polytopes and a hierarchy of combinatorial problems, *Disc. Math.* 4 (1973), 305–337.
- [14] V. Chvátal, W. Cook and M. Hartmann, On cutting-plane proofs in combinatorial optimization, *Linear Algebra Appl.* 114/115 (1989), 455–499.

- [15] J. Edmonds, Paths, trees, and flowers, *Canadian Journal of Mathematics* 17, 449–467, 1965.
- [16] J. Edmonds, Maximum matching and a polyhedron with 0-1 vertices, *Journal of Research National Bureau of Standards, Section B* 69, 73–77, 1965.
- [17] F. Eisenbrand and A.S. Schulz, Bounds on the Chvátal rank of polytopes in the 0/1-cube, *Combinatorica* 23, 245–261, 2003.
- [18] M.L. Fisher, G.L. Nemhauser and L.A. Wolsey, An analysis of approximations for maximizing submodular set functions - II. *Math. Prog. Study*, 8:73–87, 1978.
- [19] T. Fujito, A 2/3-approximation of the matroid matching problem, in “Proceedings of ISAAC 1993,” *Lecture Notes in Computer Science* v.762, 185–190.
- [20] H.N. Gabow and M. Stallmann, An augmenting path algorithm for linear matroid parity, *Combinatorica* 6, 123–150, 1986.
- [21] D. Gijswijt and G. Pap, An algorithm for weighted fractional matroid matching, TR-2008-11, Technical Report of the Egervary Research Group, Budapest.
- [22] M. Hartmann, “Cutting Plane Proofs and the Complexity of the Integer Hull,” Ph.D. Thesis, Cornell University, Ithaca, 1988.
- [23] D. Hausmann, B. Korte and T. Jenkyns. Worst case analysis of greedy type algorithms for independence systems, *Math. Prog. Study* 12 (1980), 120–131.
- [24] E. Hazan, S. Safra and O. Schwartz. On the complexity of approximating k -set packing. *Computational Complexity*, 15(1), 20–39, 2006.
- [25] C. Hurkens and A. Schrijver, On the size of systems of sets every t of which have an SDR, with an application to the worst-case ratio of heuristics for packing problems, *SIAM J. Disc. Math.* 2(1) (1989), 68–72.
- [26] P.M. Jensen and B. Korte, Complexity of matroid property algorithms. *SIAM Journal on Computing* 11, 184–190, 1982.
- [27] T.A. Jenkyns, “Matchoids: A generalization of matchings and matroids,” Ph.D. thesis, University of Waterloo, Waterloo, Ontario, 1974.
- [28] T.A. Jenkyns, The efficacy of the “greedy” algorithm, in: “Proc. of 7th SE Conf. on Comb., Graph Theory and Comp.,” 341–350, 1976.
- [29] A. Karlin, C. Mathieu and C. Thach Nguyen, Integrality gaps of linear and semi-definite programming relaxations for knapsack, manuscript, 2009.
- [30] M. Laurent, A comparison of the Sherali-Adams, Lovász-Schrijver and Lasserre relaxations for 0-1 programming. *Mathematics of Operations Research*, 28(3):470–496, 2003.
- [31] E.L. Lawler, “Combinatorial Optimization: Networks and Matroids,” Holt, Rinehart and Winston, 1976.
- [32] E.L. Lawler, Matroids with parity conditions: a new class of combinatorial optimization problems, Memo. No. ERL-M334, Electronics Research Laboratory, College of Engineering, UC Berkeley, 1971.
- [33] J. Lee, M. Sviridenko and J. Vondrák, Submodular maximization over multiple matroids via generalized exchange properties, “Proc. of APPROX,” Springer LNCS, 244–257, 2009.
- [34] J. Lee, M. Sviridenko and J. Vondrák, Matroid Matching: the Power of Local Search, IBM Research Report RC24898, 11/2009.
- [35] L. Lovász, Matroid matching and some applications, *J. Comb. Th., Ser. B* 28, 208–236, 1980.
- [36] L. Lovász, The matroid matching problem. In: L. Lovász, L., V.T. Sós (Eds.), “Algebraic Methods in Graph Theory, Vol. II,” (Colloq. Szeged, 24–31 August 1978). *Colloq., Math. Soc. Janos Bolyai* 25, North-Holland, 495–517, 1981.
- [37] L. Lovász and M.D. Plummer, “Matching Theory,” *Annals of Disc. Math.* 29, North-Holland, 1986.
- [38] C. Mathieu and A. Sinclair, Sherali-Adams relaxations of the matching polytope, in *Proc. STOC 2009*.
- [39] J. Mestre, Greedy in approximation algorithms, in: “Proc. of ESA,” 528–539, 2006.
- [40] H. Narayanan, H. Saran and V. Vazirani, Randomized Parallel Algorithms for Matroid Union and Intersection, With Applications to Arborescences and Edge-Disjoint Spanning Trees, *SIAM Journal of Computing* 23(2), 387–397, (1994).
- [41] J.B. Orlin, A fast, simpler algorithm for the matroid parity problem, In: A. Lodi, A. Panconesi and G. Rinaldi (Eds.), “Proceedings of the The 13th IPCO,” (Bertinoro, Italy, 26–28 May 2008), LNCS 5035, 2008, 240–258.
- [42] J.B. Orlin and J.H. Vande Vate, Solving the linear matroid parity problem as a sequence of matroid intersection problems,” *Mathematical Programming*, 47, 81–106, 1990.
- [43] A. Recski, “Matroid theory and its applications in electric network theory and in statics,” *Algorithms and Combinatorics* 6, Springer Verlag, Berlin, 1989.
- [44] A. Recski and J. Szabó, On the generalization of the matroid parity problem. In: A. Bondy, J. Fonlupt, J.-L. Fouquet, J.-C. Fournier, J.L. Ramírez Alfonsín (Eds.), “Graph Theory in Paris,” Trends in Mathematics, Birkhauser, Basel, 2007, 347–354.
- [45] A. Schrijver, “Theory of Linear and Integer Programming,” Wiley, 1986.
- [46] A. Schrijver, “Combinatorial Optimization: Polyhedra and Efficiency,” Springer-Verlag, Berlin, 2003.
- [47] H.D. Sherali and W.P. Adams, A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems, *SIAM J. on Disc. Math.* 3, 411–430, 1990.
- [48] M. Stallmann and H.N. Gabow, An augmenting path algorithm for the parity problem on linear matroids. In: “Proceedings of the 25th FOCS,” (Singer Island, Florida, 24–26 October 1984). IEEE, Washington, DC, 1984, 217–228.
- [49] P. Tong, E. Lawler and V. Vazirani, Solving the weighted parity problem for gammoids by reduction to graphic matching, In: “Progress in combinatorial optimization (Waterloo, Ont., 1982),” 363–374, Academic Press (Toronto, ON), 1984.
- [50] J. Vande Vate, Fractional matroid matchings, *Journal of Combinatorial Theory, Series B* 55 (1992), 133–145.
- [51] V.V. Vazirani, “Approximation Algorithms,” Springer-Verlag, Berlin, 2001.
- [52] D.J.A. Welsh, “Matroid Theory,” Academic Press, London-New York, 1976.
- [53] S.G. Williamson, “Combinatorics for computer science,” Comput. Sci. Press, Rockville, MD, 1985.